



InnoCare Pharma

Interim Results

Stock Code: 09969.HK, 688428.SH

August 24, 2026



These materials are for information purposes only and do not constitute or form part of an offer or invitation to sell or issue or the solicitation of an offer or invitation to buy or subscribe for securities of InnoCare Pharma Limited (the “Company”) or any of its holding company or subsidiaries in any jurisdiction. No part of these materials shall form the basis of or be relied upon in connection with any contract or commitment whatsoever.

The information or opinions contained in these materials has not been independently verified. No representation or warranty, whether expressed or implied, is made as to, and no reliance should be placed on, the fairness, accuracy, completeness or correctness of such information or opinions contained herein. The information and opinions contained in these materials are provided as of the date of the presentation, are subject to change without notice and will not be updated or otherwise revised to reflect any developments, which may occur after the date of the presentation. The Company, any of its affiliates, directors, supervisors, senior managers, officers, employees, advisers and their respective representatives shall not have any liability whatsoever (in negligence or otherwise) for any loss howsoever arising from or in reliance upon any information contained or presented in or derived from these materials or otherwise arising in connection with these materials.


These materials contain statements that reflect the Company’s current beliefs and expectations about the future as of the respective dates indicated herein. These forward-looking statements are based on a number of assumptions about the Company’s operations and businesses and on factors beyond the Company’s control, and are subject to significant risks and uncertainties, and, accordingly, the actual results may differ materially from these forward-looking statements. You should not place undue reliance on any of such forward-looking information. The Company assumes no obligation to update or otherwise revise these forward-looking statements for new information, events or circumstances that emerge subsequent to such dates.

Our Mission & Vision: Science Drives Innovation For The Benefit of Patients



To Become
a **Global Biopharmaceutical Leader**
that Develops and Delivers
Innovative Therapies for Patients **Worldwide**

Cancer



Autoimmune

Our Therapeutic Focus


**STRONG
FINANCIAL
PERFORMANCE
AND SIGNIFICANT
PROGRESS
ACROSS OUR
DIVERSIFIED
INNOVATION
PORTFOLIO**

FINANCIAL HIGHLIGHTS

- ❖ **2026H1: Strong drug sales growth and sustained profitability**
 - Total revenue reached **RMB 1,137.1mn**, representing a **55.5%** yoy growth
 - Drug sales achieved **RMB 918.1mn** with **43.2%** yoy growth
 - Net profit reached **RMB 239.7 million**, reflecting continued strong earnings growth
- ❖ **Cash position:** Maintained a solid cash position of **~ RMB 8.4bn by end of 2026H1**, providing ample resources to support ongoing R&D and globalization initiatives

R&D PIPELINE HIGHLIGHTS

- ❖ **1 NDA approval:** Orelabrutinib for r/r MCL in Australia
- ❖ **2 NDA submissions:** Orelabrutinib for ITP, Zurletrectinib for NTRK+ pediatric patients
- ❖ **2 Ph3 primary endpoints met:** Soficitinib for atopic dermatitis, Fadeucravacitinib for psoriasis
- ❖ **3 New Ph3 trials initiated:** Orelabrutinib for SLE, Mesutoclax + orelabrutinib for r/r MCL, Mesutoclax+AZA vs. Venetoclax+AZA in 1L AML
- ❖ **2 POC achievements:** Soficitinib for Vitiligo, B7H3-ADC
- ❖ **4 INDs Filed, with 3 Programs Advancing into Clinic:** IL-17i, VAV1 molecular glue, CDH17-ADC, PSMA/STEAP1-ADC



Hemato-Oncology

❖ Orelabrutinib


- 1L CLL/SLL NDA approval and NRDL inclusion, constantly expanding growth potential
- Overseas progress: approved for r/r MCL in Australia, approved for r/r MCL and r/r MZL in Singapore previously

❖ Tafasitamab

- Entering its first full-year of commercial sales in China for r/r DLBCL patients

❖ Mesutoclax (ICP-248)

- Combo with Orelabrutinib for **1L CLL/SLL-FDT Ph3** registrational trial - completion expected in 1H2027
- Registrational study for **BTKi treated MCL** ongoing
- Combo with Orelabrutinib for **r/r MCL Ph3** registrational trial initiated
- Combo with Orelabrutinib for **r/r MZL**: granted BTD
- **Mesutoclax + AZA vs venetoclax + AZA**: Ph3 registrational study is initiating for **elderly/unfit TN-AML**
- Global Phase 2 trial in **MDS** ongoing with promising preliminary results



Solid Tumor

❖ Zurletrectinib (ICP-723)

- Pediatric (2-<12y) NDA accepted with priority review
- 2nd-gen pan-TRK inhibitor, Dec-2025 NMPA approval (adult & ≥12y adolescent), commercial sales launched

❖ ICP-B794 (B7-H3 ADC)

- Phase I dose escalation ongoing with promising preliminary results

❖ ICP-B208 (CDH17 ADC)

- IND approved in May 2026, Phase I dose escalation ongoing

❖ ICP-B381 (PSMA/STEAP1 ADC)


- IND submitted and accepted by NMPA in Aug. 2026
- U.S. IND filing expected in Sept. 2026

Advancing a Broad **Autoimmune** Portfolio Toward Commercialization, with First-in-Class Potential in Next-Wave Assets



Late-stage Registrational Pipeline

- ❖ **Orelabrutinib**
 - **ITP**, NDA accepted in 1H2026
 - **SLE**, promising Ph2b data, Ph3 registrational trial ongoing
 - **PPMS & SPMS**, Global Ph3 trials advancing with Zenas's platform
- ❖ **Soficitinib (ICP-332)** (TYK2/JAK1)
 - **AD**: Ph3 met primary endpoints; NDA planned after safety follow-up
 - **Vitiligo**: Ph2 met primary endpoint; advancing to Ph3
 - **PN**: Global Ph2 patient enrollment ongoing
 - **CSU**: Phase 2 enrollment completed; data readout by YE2026
 - **PsO**: Phase 2 enrollment completed; data readout by YE2026
- ❖ **Fadeucravacitinib (ICP-488)** (TYK2, allosteric)
 - **PsO**: Ph3 registrational trial met its primary endpoints
 - **CLE**: Ph2 trial ongoing
 - **SS**: Ph2 trial ongoing
 - More indications under discussion



Novel early-stage assets

- ❖ **ICP-054** (IL17 small molecule)
 - Oral IL-17 inhibitor offering a differentiated alternative to injectable biologics
 - Ex-Greater China & Southeast Asia rights licensed to Zenas, with Phase I completion expected by YE2026
- ❖ **ICP-538** (VAV1 molecular glue)
 - Potential first-in-class VAV1 molecular glue degrader targeting a novel immune regulatory pathway
 - Phase I study is expected to complete by YE2026, with Phase I data expected to provide initial safety, PK/PD and target engagement insights
- ❖ Multiple programs advancing toward clinical development

Innovative Pipeline: Accelerating Portfolio Towards Value Realization



Pre-IND		Phase 1/2		Phase 3		Registration		Approved	
Degrader	Oral	Mesutoclax (ICP-248)	BCL2	Orelabrutinib	BTk	Orelabrutinib	BTk	Orelabrutinib	BTk
● Autoimmune diseases		● r/r NHL(CHN, US)		● TN MCL (Global)		● ITP (CHN)		● TN CLL/SLL (CHN)	
Biologics		● AML(Global)		● MZL confirmatory (CHN)		Zurletrectinib	NTRK	● r/r CLL/SLL (CHN)	
● Solid tumor	BsAb-ADC	● MDS (CHN, Global)		● SLE (CHN)		● NTRK fusion-positive cancers in pediatric patients (CHN)		● r/r MCL (CHN)	
● IBD	BsAb	Soficitinib (ICP-332)	TYK2/JAK1	● PPMS (Global)*				● r/r MCL (SG)	
Others	Oral	● Prurigo nodularis (Global)		● SPMS (Global)*				● r/r MCL (AU)	
● Autoimmune diseases		● Psoriasis (CHN)		Tafasitamab	CD19			● r/r MZL (CHN)	
		Fadeucravacitinib(ICP-488)	TYK2	● DLBCL (CHN)				● r/r MZL (SG)	
		● CLE (CHN)		Mesutoclax	BCL2			Tafasitamab	CD19
		● Sjögren's syndrome (CHN)		● TN CLL/SLL (CHN)	+Orela			● r/r DLBCL (CHN Mainland)	
		ICP-538	VAV1	● BTKi failure r/r MCL	Phase 2 registrational			● r/r DLBCL (GBA)	
		● Autoimmune diseases (CHN)		● r/r MCL	+Orela			● r/r DLBCL (HK)	
		ICP-054	IL-17AF*	● r/r MZL	+Orela			● r/r DLBCL (Macao)	
		● Autoimmune diseases (CHN)		● 1L AML(CHN)	+AZA vs.Ven.+AZA			● r/r DLBCL (TW)	
		ICP-B794 (ADC)	B7H3	Soficitinib (ICP-332)	TYK2/JAK1			Zurletrectinib	NTRK
		● Solid Tumors (CHN)		● Atopic Dermatitis (CHN)				● NTRK fusion-positive cancers (CHN)	
		ICP-B208(ADC)	CDH17	● Vitiligo (CHN)	Phase 2/3				
		● Solid Tumors (CHN)		● CSU (CHN)	Phase 2/3				
		ICP-B381(BsAb-ADC)	PSMA/STEAP1	ICP-488	TYK2				
		● Prostate Cancer		● Psoriasis (CHN)					

- Hemato-oncology
- Autoimmune Disease
- Solid Tumors


Total Revenue in first half 2026 achieved **55.5% yoy growth**, of which Drug Sales growing **43.2% yoy** with a Diversified Portfolio

Total Revenue

In RMB millions


BD and service

Drug sales



Drug Sales

In RMB millions




- **Total Revenue achieved 55.5% yoy growth**
 - Drug sales continued robust growth
 - BD milestone payments recognized in 2026 H1
- **Drug sales achieved 43.2% yoy growth, more commercialized products with new indications ensuring continued high growth**
 - Expanded market for Orelabrutinib after 1L CLL/SLL approval and further penetration in high-potential MZL market
 - Contributions from a diversified portfolio following the launches of Tafasitamab and Zurlitrectinib
 - Enhanced commercial execution to gain more market share

Note : The above financials are based on HKFRS (Hongkong Financial Reporting Standards)
Drug Sales include products revenue of Orelabrutinib, tafasitamab, and Zurlitrectinib

Sustainable Profitability Backed by Strong Cash Reserves

Profit/(Loss) for the Period


In RMB millions



InnoCare continued to be profitable in the first half 2026, reporting a profit of RMB 240M (Diluted EPS RMB 0.14). This achievement was primarily attributable to a strong drug revenue growth, BD milestone revenue recognized, alongside the improved cost efficiency.

R&D Expense

In RMB millions



R&D expenses increased for strategic investment for innovative technology platform, increased resources to clinical trials for our prioritized programs

Cash and related balance*

In RMB millions




Robust cash and related account balance of RMB 8.4B (~US\$1.24B) provides flexibility to expedite the clinical development and to invest in a competitive pipeline

Orelabrutinib + Tafasitamab + Zurletrectinib: Diversified Product Portfolio Driving Continued Commercial Growth



宜诺凯 Orelabrutinib **明诺凯** Tafasitamab **宜诺欣** Zurletrectinib



Orelabrutinib



- ✓ Excellent efficacy and safety profile
- ✓ **Once-daily dosing**
- ✓ **Largest patient population coverage in NHL** (CLL/SLL, MCL, MZL) among BTKi in CN
- ✓ **First and only** BTKi for the treatment of **r/r MZL**
- ✓ **Recommended in the 2026 CSCO Lymphoma Guidelines:** CLL/SLL (1L & r/r) – Grade I; MCL (1L) – Grade II, (r/r) – Grade I; MZL (r/r) – Grade I, 1L (in combination with rituximab) – Grade II; PCNSL (1L, in combination with HD-MTX and rituximab) – Grade II.

Tafasitamab



- ✓ High response rates and durable remissions
- ✓ The **first CD19-targeted antibody** approved in Greater China for **r/r DLBCL**
- ✓ **Recommended in the 2026 CSCO Lymphoma Guidelines:** r/r DLBCL – Class I; r/r FL (in combination with rituximab and lenalidomide) – Class I

Zurletrectinib



- ✓ Durable deep responses, strong CNS penetration, and favorable overall safety profile
- ✓ China's **first** domestically developed **next-generation TRK inhibitor** with tumor-agnostic potential
- ✓ Demonstrated ability to overcome resistance to first-generation TRK inhibitors

Strong commercial execution driving sustained growth

¹Indications included in NRD: adult patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) (1L CLL/SLL), adult patients with mantle cell lymphoma who have received at least one prior therapy (r/r MCL), and adult patients with marginal zone lymphoma who have received at least one prior therapy (r/r MZL)

A person wearing a blue cleanroom suit, hood, and mask is reviewing a large sheet of paper in a complex industrial setting with white pipes and machinery. The background is slightly blurred, emphasizing the person and their work.

A Leading Hemato- oncology Franchise

Hemato-oncology: Leading Commercializing Proven Portfolio and Advancing Next-Generation Therapies for Broader Markets




Assets	Target	Indication	Clinical Trial	Registration	Market
Orelabrutinib	BTk	r/r CLL/SLL			★ CHN
		r/r MCL			★ CHN,SG,AU
		r/r MZL			★ CHN, SG
		1L CLL/SLL			★ CHN
		1L MCL	Global Ph3 ongoing	🎯	
		MZL Confirmatory Trial	Ph3 ongoing	🎯	
Tafasitamab	CD19	r/r DLBCL			★ CHN, HK, MC, TW
		DLBCL Confirmatory Trial	Ph3 ongoing	🎯	
Mesutoclax (ICP-248)	BCL2	1L CLL/SLL	+Orela, Ph3, patient enrollment completed	🎯	
		r/r MCL (BTKi treated)	Ph2 registrational trial	🎯	
		r/r MCL	+Orela, Ph3 registrational trial	🎯	
		r/r MZL	+Orela, Ph3 registrational trial	🎯	
		1L AML	+AZA vs Ven+AZA, Ph3 registrational trial	🎯	
		1L MDS	Dose escalating ongoing in CHN & global		

★ Market

🎯 Registration trial

Tafasitamab: Best Medicine Potential to Deliver Unique Clinical Value for DLBCL Patients




Comparison of Selected Novel Therapy in r/r DLBCL

Company	Target	Therapy	Phase	ORR (%)	CR (%)	mDOR (m)	mPFS (m)	mOS (m)
Incyte/InnoCare	CD19	Tafasitamab + Lenalidomide	Approved ex-China	57.5	40	43.9	11.6	33.5
ADC Therapeutics	CD19 ADC	Loncastuximab tesirine	Approved ex-China	48.3	24.1	10.25	4.93	9.92
Roche	CD79b ADC	Polatuzumab vedotin + BR	Approved	42	23	12.6	9.5	12.4
Roche	CD20/CD3	Glofitamab	BLA	52	39	10.4	3.8	11.5
Amgen/Beigene	CD19/CD3	Blinatumomab	II	43	19	11.6	3.7	5.0
Regeneron/Zai Lab	CD20/CD3	Mosunetuzumab	II	33	21	N/A	N/A	N/A
AbbVie	BCL-2	Venetoclax+R+Pola	II	65	31	5.8	4.4	11

Non-head-to-head comparison

Mesutoclax (ICP-248): A Novel BCL-2 Inhibitor with Great Clinical Advantages



Venetoclax Pharmacological Properties

- M27, a major metabolite of Venetoclax, shows ~80% AUC of the parent drug within 24 h
- M27 has no pharmacological activity but has hematological toxicity*
- Significant inhibition of CYP2C8 and CYP2C9 by Venetoclax and M27 with $\text{IC}_{50} \leq 0.82 \mu\text{M}$
- Significant inhibition of P-gp and BCRP by Venetoclax and M27 with $\text{IC}_{50} \leq 1.48 \mu\text{M}$

Advantages of Mesutoclax



Eliminated major metabolite



Significantly higher exposure



Reduced hematological toxicity



Reduced DDI risks




Excellent efficacy & safety profile

* Venetoclax FDA non-clinical toxicology review

Favorable PK Profiles Compared with Venetoclax in Clinical Studies

Comparison of PK Profiles
ICP-248 (100/125 mg, QD) vs Venetoclax (400 mg, QD)



✓ PK exposure of ICP-248 at **125 mg QD** is **3 times** of Venetoclax at **400 mg QD**

Mesutoclax (ICP-248): Strengthening Our BTK + BCL-2 Franchise to Drive Leadership in NHL



1L CLL/SLL-FDT

Combo with Orelabrutinib, Ph3 registrational trial

	Orela+Mesutoclax	Ibru + Ven ¹	Acala + Ven ²
Sample Size	42	106	291
ORR	100%	86.8%	92.8%
CRR	52.4%*	36.7%	NA
uMRD	65%** W36	45.3% EOT+3	34.4% EOT
TLS	0	0	0.3%

Cutoff date: 2026-01-26

* Complete remission in target lesion at RP3D per image

** MRD checkpoint at 36th week of combo treatment, in patients achieved CR

BTKi-treated MCL

Registrational trial, First BCL-2 inhibitor in China to receive Breakthrough Therapy Designation

	Mesutoclax	Venetoclax ^{3,4}	Pirtobrutinib ⁵
	BTKi+, N=25	BTKi+, N=17	cBTKi* Pretreated MCL N=90
ORR	84%	53%	57.8%
CRR	36%	18%	20.0%

Cutoff date: 2025-07-10

* cBTKi: covalent Bruton tyrosine kinase inhibitor

r/r MZL

Combo with Orelabrutinib, Ph3 registrational trial

- Early clinical data showed 100% ORR and 50% CRR in evaluable r/r MZL patients
- BTD granted by CDE for Orelabrutinib + Mesutoclax in r/r MZL after ≥1 prior therapy
- Ph3 IND application submitted

r/r MCL

Combo with Orelabrutinib, Ph3 registrational trial

- Mesutoclax + Orelabrutinib demonstrated encouraging activity in r/r MCL, achieving 100% ORR and 100% CRR among evaluable patients
- Ph3 study of Mesutoclax + Orelabrutinib in r/r MCL approved to initiate in China

Mesutoclax (ICP-248): Advancing AML and MDS Programs with Significant Global Market Potential



1L AML:
Ph3 Mesutoclax+AZA vs. Venetoclax+AZA in China
Global dose expansion

MDS: Global Ph2 study ongoing

1L AML

	Mesutoclax	Venetoclax ¹	Lisaftoclax ²	Sonrotoclax ³
	N=44	N=286	N=39	N=79
CRR	81.8%	66.4%	51.3%	67.1%
uMRD*	86.5%	23.5%	NA	52.8%
SAE	20.5%	83%	43.3%	77.2%
6-month OS	90.5%	71.9%	/	/

Cutoff date: 13th Apr. 2026


MDS

Response per IWG 2006	Efficacy Evaluable N=10
ORR	100%
CR	40%
mCR	60%

Cutoff date: 20th Apr. 2026

1. N Engl J Med 2020;383:617-29.
2. 2024 ASCO
3. 2025. EHA
4. Nova One Advisor, Insight Code: 8817
Note: *Calculate in patients with composite complete remission

Ph3 Randomized, Open-Label, Multicenter Study of Mesutoclax + AZA vs. Venetoclax + AZA in 1L AML



Mesutoclast (ICP-248): Best-in-Class Potential and Significant Market Opportunity

1L CLL/SLL Fix-duration Treatment

- The estimated CLL/SLL patient population in China is approximately 29,000¹.
- The market is currently valued in the **billions of RMB** and is expected to expand following the approval of **fixed-duration therapies**.

r/r MCL

- BTK inhibitors are broadly established in treating **MCL**².
- With growing **BTKi-resistance**, the unmet need for subsequent therapies is substantial.

r/r MZL

- MZL is the second-largest NHL indication, representing a significant market opportunity.
- BCL-2 combination provides a significant opportunity to **expand patient penetration** and **reinforce market leadership**.

AML

- Global **new AML** cases are projected to increase from ~103K in 2018 to ~115K by 2028³.
- The global AML market was estimated at **US\$3.7 billion** in 2024 and is expected to grow to **US\$8 billion** by 2034³.

MDS


- Global **MDS** patient population: approximately 500K⁴.
- The global myelodysplastic syndrome drugs market size was valued at **US\$4.6 billion** in 2024 and is anticipated to reach around **US\$11 billion** by 2034⁵.

Addressable Market Potential: ~ US\$20 billion

Well Positioned Portfolio in Autoimmune Diseases



Multiple Assets with Large Indications Progressed to Phase 3 Trials



PPMS: Primary Progressive Multiple Sclerosis
 SPMS: Secondary Progressive Multiple Sclerosis
 ITP: Idiopathic Thrombocytopenic Purpura
 CSU: Chronic Spontaneous Urticaria
 SS: Sjögren's Syndrome
 SLE: Systemic Lupus Erythematosus
 AD: Atopic Dermatitis
 PN: Prurigo Nodularis
 CLE: Cutaneous Lupus Erythematosus

Orelabrutinib: Enormous Potential for Treating Autoimmune Diseases



MS¹

- **PPMS: Global Ph3 ongoing**
- **SPMS: Global Ph3 ongoing**
- Best-in-class potential
- Current SPMS and PPMS commercial opportunity in the U.S. alone projected to be **>\$12B²**, expected to grow significantly with approval of effective therapies that impact disease progression

SPMS & PPMS could represent >40% of all MS diagnoses

ITP¹

- Ph3 registrational trial for the treatment of ITP is underway in China, with **NDA submitted in 2026H1**
- BTKi treatment for autoimmune diseases is just around the corner

Over 200,000 new patients globally each year

SLE¹

- The **world's first and only** BTKi demonstrating efficacy in Ph2 trial
- **Phase 2b** clinical trial **met primary endpoints**
- **Phase 3 clinical trial ongoing**

~8 million patients worldwide

Orelabrutinib in ITP: Large Market Opportunity Approaching NDA

Disease & Patient Population

- ITP (Immune Thrombocytopenia) is a chronic autoimmune bleeding disorder with significant relapse rates after first-line therapy.
- ~300,000 chronic patients in China
- ~60,000 new cases annually

Current Treatment Gaps

Current Therapy	Limitations
Steroids & IVIG	Short-term benefit, significant side effects
TPO-RA	Risk of thrombotic events, decreased efficacy with prolonged treatment
Others	Lack of durable, safe oral options

Orelabrutinib's Advantage

Inhibits **abnormal B-cell** activation & **autoantibody production** with wider safety margin and convenient oral dosing

Market Potential

China's large ITP patient base and growing diagnosis rate create a significant market opportunity worth hundreds of **millions USD**

Key Milestones

2026 H1: NDA submitted and accepted by CDE

Poised to address the significant unmet needs in ITP – strong potential to become the next growth driver.

Orelabrutinib in SLE: Global First-in-Class BTK Inhibitor with Large Market Opportunity

First-in-Class potential, unlocking a multi-billion-dollar market opportunity.

- Ph3 registrational trial ongoing

SLE

- SLE is a chronic autoimmune disease affecting multiple organs.
- ~8 million patients globally; ~1 million in China¹.
- Most common in young and middle-aged women; chronic management needed for years or decades.

Orelabrutinib's Advantage

- Selective BTK inhibition to suppress B-cell activation and autoantibody production
- Oral dosing with favorable safety and tolerability
- Potential to become the first-in-class oral BTK inhibitor for SLE, offering improved convenience and disease control


Current Treatment Gaps

Current Therapy	Limitations
Corticosteroids & Immunosuppressants	Significant toxicity, poor long-term safety, frequent relapse after dose reduction
Biologics	High cost, IV or SC administration, partial response in many patients
Others	Lack of durable, safe oral options

Market Potential

- Large and underserved SLE population with increasing diagnosis rates
- Biologics market for SLE already exceeds **US\$3 billion globally**, expected to grow rapidly with more accessible oral options


ICP-332, ICP-488: TYK2 Inhibitors with Different Selectivity Profiles



Active site binding

JH2
(pseudokinase domain)

JH1
(kinase domain)



Allosteric site binding

Blocking the ATP binding site

↓
Inactive state




Inhibitor	IC ₅₀ (nM)	IC ₅₀ (nM) @1 mM ATP			
	TYK2 JH2	TYK2 JH1	JAK1	JAK2	JAK3
ICP-332	2319	0.5	19	191	930
ICP-488	5	>10,000			

Soficitinib (ICP-332): Shows Outstanding EASI 75 Improvement and Rapid Itch Reduction in Ph2 Atopic Dermatitis (AD) Trial


Soficitinib (ICP-332) Ph2 AD

Phase 2 data indicates that soficitinib demonstrates significant efficacy in treating AD, showing the best efficacy on EASI 75 (placebo-adjusted) compared to several other innovative drugs



Not a head-to-head comparison

Pruritus Numerical Rating Scale



Source: 1,2,3,4,5,6: data from ClinicalTrials.gov <https://www.clinicaltrials.gov/>

7. DUPIXENT® (dupilumab) injection label.

8. A. Wollenberg, et al. Br J Dermatol 2021; 184:386–387 DOI 10.1111/bjd.19574.


9. Silverberg JI, et al. N Engl J Med . 2023 Mar 23;388(12):1080-1091. doi:

Soficitinib (ICP-332): Showed Excellent Efficacy in Ph3 AD Study with All Primary Endpoint Met

A phase 3 randomized, double-blind, placebo-controlled study to evaluate the efficacy and safety of oral ICP-332 in subjects with moderate to severe atopic dermatitis (NCT06775860)

▪ Patient Population:

- EASI ≥ 16
- $\geq 10\%$ BSA
- vIGA-AD ≥ 3
- Baseline average WP-NRS ≥ 4



▪ CO-Primary Endpoints:

- EASI 75 at W16
- vIGA 0/1 and ≥ 2 points reduction at W16

▪ Secondary Endpoints:

- EASI 50 /75 /90 /100
- vIGA 0/1 and ≥ 2 points reduction
- NRS 0/1
- Improvement of Worst Pruritus NRS ≥ 4
- Time to achieve improvement of Worst Pruritus NRS ≥ 4


- 16-week efficacy primary endpoints successfully met, marking a key clinical milestone
- 52-week safety follow-up ongoing, with NDA submission planned upon completion

Soficitinib (ICP-332): Positive Ph2 Vitiligo Results Pave the Way for Ph3

A phase 2/3 randomized, double-blind, placebo-controlled, parallel group, adaptive design, multi-center study to evaluate the efficacy and safety of ICP-332 in subjects with non-segmental vitiligo (NCT07047612)

■ Patient Population:

- Non-segmental vitiligo subjects



■ Primary Endpoints:

- F-VASI change from baseline at W24


■ Secondary Endpoints:

- F-VASI 50/75/90/100
- T-VASI 50/75/90/100
- PGIS-F
- PGIS-V
- VNS 4 / 5
- HADS
- VitiQoL
- Safety
- PK and biomarker

- Ph2 vitiligo study met its primary endpoint, demonstrating positive clinical efficacy
- Ph3 initiation underway, advancing the program toward pivotal development

Soficitinib (ICP-332) Ph2 Vitiligo Study: Primary End Point Met with Excellent Efficacy on F-VASI Improvement from Baseline at W24

F-VASI Improvement from Baseline at W24



Soficitinib (ICP-332): Enormous Potential for Treating Inflammatory Skin Diseases


 ICP-332 Atopic Dermatitis (AD)	 ICP-332 Vitiligo	 ICP-332 CSU	 ICP-332 Psoriasis	 ICP-332 Prurigo Nodularis (PN)
• Ph3 met primary endpoints; NDA submission planned in 2027H1. • TYK2/JAK1 inhibitor blocks key cytokine signaling pathways: IL-4, IL-13, IL-31, and TSLP, suppressing Th2-driven inflammation and alleviating atopic dermatitis symptoms. • Global drug-market for AD: estimated at US\$18 billion in 2024, projected to reach ~ US\$30 billion by 2030¹.	• Ph2 positive; Ph3 initiation underway. • TYK2/JAK1 inhibitor blocks IFN-γ and IL-15–mediated JAK-STAT signaling, suppressing T-cell attacks on melanocytes and promoting repigmentation. • The global vitiligo treatment market size was valued at US\$2 billion in 2024 and is projected to reach US\$3 billion by 2032³.	• Ph2/3 ongoing; Ph2 enrollment completed, with data readout expected by YE2026. • TYK2/JAK1 inhibitor blocks cytokine signaling pathways: IL-4, IL-13, and IL-31 that drive mast cell activation and inflammation, reducing itch and wheal formation in CSU. • The global CSU treatment market has reached to US\$2 billion in 2024 and expected to grow to US\$3 billion in 2029⁵.	• Ph2 enrollment completed; data readout expected by YE2026. • Dual TYK2/JAK1 pathway modulation suppresses IL-23/Th17 signaling and multiple pro-inflammatory cytokines, reducing inflammation and normalizing keratinocyte hyperproliferation. • Global psoriasis treatment market: ~US\$27 billion in 2024, projected to reach ~US\$58 billion by 2032¹⁷.	• Global Ph2 ongoing. • TYK2/JAK1 inhibitor blocks cytokine signaling pathways: IL-4, IL-13, and IL-31, reducing neurogenic pruritus and alleviating PN symptoms. • The global PN market was valued at US\$2 billion in 2024 and is expected to grow to US\$3 billion in 2034⁹.
~200 million patients worldwide²	~70 million patients worldwide⁴	~50 million patients worldwide⁶	240 million patients worldwide⁸	~10 million patients globally¹⁰

★ Met primary endpoint(s).


☆ Data coming in the next few months.

Fadeucravacitinib (ICP-488): Shows Strong Efficacy in Ph2 Plaque Psoriasis

Patients achieving PASI 75 at Week 12 (FAS)



PASI 75 Response Rate by visit (FAS)



All randomized subjects were included in the FAS analysis. p values from a Cochran-Mantel-Haenszel test, with prior biologic treatment included as a stratification factor, comparing the proportion of patients in the treatment group versus placebo.


PASI, Psoriasis Area and Severity Index; QD, once daily; NRI, non-responder imputation

Fadeucravacitinib (ICP-488): Showed Excellent Efficacy in Ph3 Psoriasis Study with All Primary Endpoint Met

A phase 3 randomized, double-blind, placebo-controlled study to evaluate the efficacy and safety of oral ICP-488 in subjects with moderate to severe plaque psoriasis (NCT06842199)

▪ Patient Population:

- PASI $\geq$ 12
- BSA $\geq$ 10%
- sPGA $\geq$ 3



▪ CO-Primary Endpoints:

- PASI 75 at W16
- sPGA 0/1 and $\geq$ 2 points reduction at W16

▪ Secondary Endpoints:

- PASI 50 /75 /90 /100
- sPGA 0/1

- 16-week efficacy primary endpoints successfully met, marking a key clinical milestone
- 52-week safety follow-up ongoing

Fadeucravacitinib (ICP-488): Leveraging Strong Efficacy and Safety Profile, Unlocking Broader Market Potential



ICP-488 Psoriasis

- **Ph3 met primary endpoints.**
- TYK2 allosteric inhibitor with potential to achieve best-in-class efficacy and safety.
- Global psoriasis treatment market: ~US\$27 billion in 2024, projected to reach ~US\$58 billion by 2032¹.

240 million patients worldwide²

ICP-488 CLE

- **Ph2 ongoing.**
- TYK2 allosteric inhibitor with potential to address high unmet need in CLE, targeting the Type I interferon pathway central to lupus pathogenesis.
- Global market estimated at ~US\$2.9B in 2024, projected to reach ~US\$7.9B by 2032³.

~800,000 diagnosed patients across major markets⁴

ICP-488 Sjögren's syndrome

- **Ph2 ongoing.**
- TYK2 allosteric inhibitor modulate type I interferon and interleukin signaling pathways, underlying immune dysregulation of Sjögren's syndrome.
- Global Sjögren's syndrome market was ~USD 3.02 billion in 2024, projected to reach ~USD 5.5 billion by 2035⁵.

3.3 million diagnosed cases across major markets⁶


ICP-488 Broad immunology platform

- **SLE:** ~8 million patients globally; ~1 million in China⁷
- **IBD:** ~US\$23.6B in 2024, projected to reach ~US\$35.7B by 2032⁸
- **Psoriatic arthritis:** ~US\$11B market growing to >US\$20B by 2030⁹

Target indications represent a combined global market opportunity of >US\$150B¹⁰

★ Met primary endpoint(s).

1. Fortune Business Insights; 2. International Federation of Psoriasis Associations (IFPA); 3,8. Data Bridge Market Research; 4. <https://www.openpr.com/news/4176533/cutaneous-lupus-erythematosus-market-epidemiology>
5. wiseguyreports; 6. delveinsight; 7. doi: 10.2478/rir-2022-0006; 9. Grand View Research; 10. Alumis




- ❑ VAV1 is a promising target involved in both T-cell and B-cell pathways that could be used for the treatment of various autoimmune diseases
- ❑ It's proposed that VAV1 targeted therapy could provide much better efficacy in some of the hard-to-treat autoimmune diseases
- ❑ ICP-538: a potent and selective VAV1 degrader. **Phase I ongoing** with SAD/MAD escalation cohorts underway

ICP-538: A Potent CRBN-Mediated VAV1 Molecular Glue Degradader with Strong Anti-Inflammatory Efficacy


Dose-response VAV1 degradation

VAV1 degradation dynamics in Jurkat cells




Rapid and deep VAV1 degradation

VAV1 degradation dynamics in Jurkat cells



Note: VAV1 degradation was tested in Jurkat cells.


ICP-538 Inhibits Rat CIA Progression




* CIA: collagen-induced arthritis

IL-17: A Valid Autoimmune Disease Target

Mechanism of Action of IL-17 Inhibitor



Target Indications




ICP-054: A Novel Small-Molecule IL-17 Inhibitor for Autoimmune Diseases with Potent Activity Against Both IL-17AA and IL-17AF

ICP-054 is potent against both IL-17AA and AF

ICP-054


- Potent against IL-17AA & AF
- Excellent PK
- **Phase I ongoing** with SAD/MAD escalation cohorts underway.



IL-17 modulator (PPI)

CMPD	Binding IC ₅₀ (nM)	IL-17 Signaling IC ₅₀ (nM)@HEK-Blue		IL-17 Signaling IC ₅₀ (nM)@HEK-Blue Th17 Supernatant
		IL-17AA	IL-17AF	
ICP-054	2.6 ± 1.7	6.5 ± 3.1	18.4 ± 16.2	2.8 ± 3.4

In vivo efficacy in rat CIA model



Innovative Therapies for Comprehensive Coverage of Autoimmune Diseases

Orelabrutinib (BTKi)

Soficitinib (ICP-332) (TYK2/JAK1i)

Fadeucravacitinib (ICP-488) (TYK2i)

ICP-538 (VAV1 molecular glue)

ICP-054 (IL-17 small molecule)


Project 42 (molecular glue degrader)

Project 40 (bi-specific antibody)

Project 43 (PROTAC)

Project 44 (small molecule)

■ Clinical
■ Pre-clinical



■ InnoCare current coverage

LN: Lupus Nephritis
MN: membranous nephropathy
UC: Ulcerative Colitis
CD: Crohn disease
CSU: Chronic Spontaneous Urticaria

AA: Aplastic anemia
AIHA: Autoimmune hemolytic anemia
NMO: Neuromyelitis optica
MG: Myasthenia gravis
CLE: Cutaneous Lupus Erythematosus

MoG-EM: MOG antibody-associated encephalomyelitis
SS: Sjögren syndrome
RA: Rheumatoid Arthritis
IgG4 RD: IgG4 related disease

A close-up photograph of a person in a white lab coat and white gloves using a pipette to transfer liquid into a small vial. The person is wearing safety glasses. The background is a blurred laboratory setting. The text "Innovative Solid Tumor Assets" is overlaid on the left side of the image.


Innovative Solid Tumor Assets

Zurletrectinib (ICP-723): 2nd Generation TRKi for NTRK Tumors




- 2nd Generation TRKi for Tumors with NTRK Gene Abnormalities – NDA for **adults and adolescents** patients, **approved in Dec. 2025**
- Registration trial for NTRK gene abnormalities **in adults and adolescents**
 - ✓ **ORR: 89.1% (95% CI: 77.8, 95.9)**
 - ✓ Long duration of response (longest beyond 36 months)
 - ✓ Efficacious in TRKi-resistant patients
- **NDA** submission for pediatric patients submitted in **2Q2026**
 - ✓ IRC-assessed **ORR: 100% (95% CI: 69.2, 100.0)**, with **10% CR** and **90% PR** as of July 11, 2026

Significant and durable efficacy observed across diverse tumor types in adult patients



Data cut-off: 2024-11-23

Design & Advantage of InnoCare's Proprietary ADC Platform



Novel Connector

- Irreversible connector
- Prevents thiol exchange

Hydrophilic Linker


- Allows high DAR
- Improves stability

Effective Payload



- Potent
- Bystander effect
- Tumor-specific release
- Rapid clearance

CDH17: A Potent Target for the Treatment of GI Cancers

Normal cells: CDH17 is hidden in tight junctions



Cancer cells: redistribution of CDH17, and making it accessible



CDH17



ICP-B208




Payload




TOPO1

ICP-B208 (CDH17 ADC): Demonstrates Robust Anti-tumor Activity Even in CDH17-low Tumors

SUN-16 (CDH17-high gastric cancer)




Gp2d (CDH17-low colorectal cancer)




- ✓ ICP-B208 is a highly potent CDH17 ADC with robust anti-tumor activity even in CDH17-low tumors
- ICP-B208 Phase I trial on-going for the treatment of gastrointestinal tumors

STEAP1/PSMA ADC: Dual-Targeting Design Expands Tumor Coverage and Addresses Antigen Heterogeneity


Heterogeneity in PSMA and STEAP1 Expression



Tumor Relapse Caused by Antigen Loss




Dual Target Binding of PSMA/STEAP1 ADC




- ✓ Potential for enhanced activity when both antigens are presented
- ✓ Good efficacy even at low target expression levels
- ✓ **Overcoming drug resistance** mediated by loss of either antigen
- ✓ IND submitted and accepted by CDE
- ✓ U.S. IND filing expected in Sept. 2026

ICP-B381: Demonstrates Robust and Dose-Dependent *In Vivo* Antitumor Activity

Robust *in vivo* antitumor activity



Dose-dependent antitumor activity



- Irreversible linker enabling high **DAR (~8)**
- Rapid payload clearance supporting a wide **therapeutic window (>250×)**
- Potent payload with activity against **large tumors** and **upon rechallenge**

- **ICP-B794 (B7H3 Target ADC)**
 - ✓ Ph1 ongoing in advanced solid tumors
 - ✓ Dose level 5 completed and dose level 6 initiated
- **ICP-B208 (CDH17 Target ADC)**
 - ✓ Ph1 trial on-going for the treatment of gastrointestinal tumors
- **ICP-B381 (STEAP1/PSMA BsAb ADC)**
 - ✓ IND submitted and accepted by CDE for the treatment of prostate cancer
 - ✓ U.S. IND filing expected in Sept. 2026

Key Upcoming Events

	Assets	Milestones
 Hemato-oncology	Orelabrutinib & Mesutoclax (ICP-248)	- Ph3 registrational trial of Orelabrutinib + Mesutoclax in 1L CLL/SLL-FDT data readout & NDA submission
		- Completion of the registration trial for BTKi-treated MCL & NDA submission
		- Ph3 initiation for r/r MCL & r/r MZL , Mesutoclax + Orelabrutinib
		- Ph3 initiation for 1L AML , Mesutoclax + AZA vs venetoclax + AZA
		- MDS : Global Phase II completion, advancing toward Phase III
 Autoimmune Diseases	Orelabrutinib	- ITP NDA Approval
		- Accelerate patient enrollment of SLE Ph3 registration trial
		- Zenas-partnered PPMS/SPMS global Ph3 programs: rapid advancement and accelerated execution
	Soficitinib (ICP-332)	- Ph3 AD , safety follow-up completion and NDA submission
		- Ph2/3 vitiligo , Ph3 initiation
		- Ph2/3 CSU & Ph2 psoriasis , Ph2 portion data readout , Ph3 initiation
	Fadeucravacitinib (ICP-488)	- Finish Global Ph2 trial in PN
		- Ph3 psoriasis , safety follow-up completion
		- Ph2 trial in CLE & SS , accelerate patient enrollment
ICP-538 (VAV1)	- Explore new indications	
	- Completion of Ph1 & data readout	
 Solid Tumor	ICP-054 (IL-17i)	- Completion of Ph1 & data readout
	ICP-B794 (B7H3 ADC)	- Dose expansion completed; recommended dose established
	ICP-B208 (CDH17 ADC)	- Dose escalation data readout and dose expansion
 Solid Tumor	ICP-B381(STEAP1/PSMA ADC)	- IND approval and FPI in China & US
	Sustained Profitability	
Sustain profit-positive performance throughout 2026		



INNOCARE

*Thank you for your
attention !*